

Therya, 2018, Vol. 9 (3): 237-246 DOI: 10.12933/therya-18-605 ISSN 2007-3364


**Predictive species distribution model of two endemic kangaroo rats from Mexico: *Dipodomys ornatus* and *D. phillipsii* (Rodentia: Heteromyidae)**


Félix J. Flores-Zamarripa<sup>1</sup> and Jesús a. Fernández<sup>1, 2\*</sup>

<sup>1</sup> Posgrado en Recursos Naturales, Facultad de Zootecnia y Ecología, Universidad Autónoma de Chihuahua. Periférico Francisco R. Almada Km 1, Zootecnia CP. 31415, Chihuahua. Chihuahua, México. Email: [felflozam@hotmail.com](mailto:felflozam@hotmail.com) (FJFZ).


<sup>2</sup> Departamento de Recursos Naturales, Facultad de Zootecnia y Ecología, Universidad Autónoma de Chihuahua. Periférico R. Almada Km 1, Zootecnia CP. 31415, Chihuahua. Chihuahua, México. Email: [afernandezf@uach.mx](mailto:afernandezf@uach.mx) (JAF).

\* Corresponding author





Map of the predictive distribution model for *Dipodomys ornatus* in Mexico



Map of the predictive distribution model for *Dipodomys phillipsii* in Mexico